

ARMY RESEARCH LABORATORY



# Chebyshev Polynomial Fit for Terrain Elevation

by Richard B. Loucks

ARL-TN-86

December 1996

19961220 082

Approved for public release; distribution unlimited.

DTIC QUALITY INSPECTED 1

# Army Research Laboratory

Adelphi, MD 20783-1197

---

ARL-TN-86

December 1996

---

## Chebyshev Polynomial Fit for Terrain Elevation

Richard B. Loucks

AMSRL-IS-EE, ARL

---

## Abstract

---

There is currently a desire to use Chebyshev polynomials to fit terrain elevation data. Such a fit would create a surface function that exactly fits the known elevations, and would describe an elevation at any point on that surface. This note questions the appropriateness of using Chebyshev polynomials for this purpose, as opposed to linear interpolation or use of a cubic spline. A set of elevations in one direction is used to illustrate a point that large transitions in elevation influence the coefficients in the polynomial fit and contribute spectral energy to points far from the transition area. It argues that a linear interpolation process, or a cubic spline interpolation, would be more appropriate.

## Contents

1. Introduction .....	1
2. Use of a Chebyshev Polynomial Curve Fit .....	2
3. Cubic Spline Interpolation .....	3
4. Conclusions .....	4
References .....	4
Distribution .....	5
Report Documentation Page .....	7

## Figures

1. Elevation data along an arbitrary isolatitude .....	2
2. Chebyshev polynomial fit to terrain data .....	3
3. Cubic spline fit to terrain data .....	4

# 1. Introduction

There is a concern whether a Chebyshev polynomial curve fit to elevation data is an appropriate method for interpolating the elevation values for points between those that are known, such as those given by the Defense Mapping Agency Digital Terrain Elevation Data, level 1 coverage. In brief, a Chebyshev polynomial is defined as (Hamming, 1986; Press et al, 1994)

$$T_n(x) = \cos(n \cos^{-1}(x))$$

or

$$T_0(x) = 1$$

$$T_1(x) = x$$

$$T_2(x) = 2x^2 - 1$$

$$T_3(x) = 4x^3 - 3x$$

...

$$T_n(x) = 2xT_{n-1}(x) - T_{n-2}(x) \quad n \geq 2$$

A curve fit using Chebyshev polynomials takes on the form

$$f(x) = \left[ \sum_{k=1}^N c_k T_{k-1}(x) \right] - \frac{1}{2} c_1$$

where the Chebyshev polynomial coefficients,  $c_k$ , are determined from

$$c_k = \frac{2}{N} \sum_{n=1}^N f(x_n) T_{k-1}(x_n)$$

where

$N$  = total number of data points to fit,

$f(x)$  = exact function described by a Chebyshev polynomial curve fit,

$f(x_n)$  = data point at location  $n$ , and

$c_k$  = Chebyshev polynomial coefficient for polynomial  $k - 1$ .

By taking advantage of the polynomial's orthogonality, we can determine the coefficients in a simpler manner by substituting for  $x_n$  with

$$x = \cos\left(\frac{\pi(k - \frac{1}{2})}{n}\right)$$

which results in

$$c_k = \frac{2}{N} \sum_{n=1}^N f\left[\cos\left(\frac{\pi(n - \frac{1}{2})}{N}\right)\right] \cos\left(\frac{\pi(k - 1)(n - \frac{1}{2})}{N}\right)$$


This last relationship is important because it gives the Chebyshev polynomial a Fourier transform quality. In essence, the coefficients are the product of a discrete Fourier transform of the elevation data. This will later give rise to problems of anti-aliasing when the curve fit is created.

## 2. Use of a Chebyshev Polynomial Curve Fit

The rationale for using a Chebyshev polynomial to fit a curve is that it will exactly fit all points considered in the fit. This type of fit uses the spectral information of the known points to determine the behavior of a function in between and over all known points. The Chebyshev polynomial is also very convenient to use since the polynomial fit at an arbitrary point need not be determined from the entire set of coefficients. There is a point in the summation where additional terms are negligible. Often times, only about 20 percent of the terms are needed to converge. In the terrain data set, it is quite easy to produce a matrix of coefficients that would represent a Chebyshev "surface." An elevation value would be simply calculated from this array of coefficients. Unfortunately, there are some problems.

Terrain data need not be continuous. Cliffs, overhangs, river gorges, canyons, mesas, or essentially any rapid transition in elevation along an isolatitude or isolongitude would present a problem in the Chebyshev fit. Remember that the Chebyshev coefficients are very similar to the coefficients generated by a Fourier transform. Consider a set of discrete points with somewhat abrupt changes in value, such as depicted in figure 1. Straight lines are drawn between the points to illustrate the effect of a linear interpolation. As is easily seen, there are some large transitions, with the largest occurring between points 1 and 2. A rough estimate of the slope is somewhat steep at  $\Delta y / \Delta x = -0.46$ . These points are not an unreasonable expectation of hilly terrain leading to a valley. In fact, they would represent a likely cross-section of an avenue for ground troop movement or encampment.

Figure 1. Elevation data along an arbitrary isolatitude.



A Chebyshev polynomial fit results in a curve that is shown in figure 2. One can immediately see the effect of anti-aliasing in the first 1000 m of the terrain profile, which is brought on by the highly transient regions contributing to the value (or energy) of the high wave number terms. These terms are later reflected in the oscillations near the base of the hill as continuing, yet decaying, in amplitude. In fact, everywhere in this section of terrain, the Chebyshev fit has introduced high spectral content in the fit. This is the occurrence in the Fourier transform of a discontinuous function. This is typically not true in the termination of hills into a valley.

### 3. Cubic Spline Interpolation

So the question becomes, is it still more reasonable to fit terrain elevation with a Chebyshev polynomial rather than use a linear interpolation, or is there something better? Perhaps the use of a cubic spline evaluated from a few surrounding points would be a better way to account for local trends, still gain a nonlinear estimate of the terrain elevation, and yet not have effects from terrain that is quite distant from the area of interest. The cubic spline presents a compromise between the linear interpolation and a Chebyshev polynomial fit. The changes between points are represented by a rounded transition to better represent hills and bluffs. Although precipitous drops are still smoothed and remain underrepresented in the terrain fit, the results are probably better than a simple linear interpolation. Looking at the same terrain data, but using a cubic spline to fit, we see in figure 3 that these two criteria are met; namely, only local effects are considered, and the curvature (in this case, based on the calculated second derivative) is used to estimate the deviation from linearity.

A cubic spline interpolation is probably better for estimating elevations, especially in the case of multiple interpolations between actual data points (desired resolution greater than actual grid spacing). A cubic spline would resolve a smooth transition about the data point. If elevation across several points is needed (desired resolution less than actual grid spacing), linear interpolation is most likely adequate, since fine terrain features are filtered out by the larger grid spacing, and average values representing the pre-

Figure 2. Chebyshev polynomial fit to terrain data.





Figure 3. Cubic spline fit to terrain data.



dominant terrain of the area are more reasonable for a fluid dynamic computation boundary. The computational cost of evaluating the Chebyshev polynomial surface, and then extracting the interpolated values, is much more costly than performing a cubic spline interpolation, as the cubic spline interpolation is more costly than linear interpolation.

## 4. Conclusion

By this assessment, the cubic spline interpolation is beneficial only if grid spacing less than twice the elevation data grid spacing is needed. Error associated with linear interpolation is probably of the same order as the error associated with a cubic spline for larger grid spacing. A Chebyshev polynomial fit near a high relief area would result in excessive elevation oscillations that may not be realistic. It is recommended that a cubic spline interpolation process be used for fine grid interpolation and that linear interpolation be used for coarse grid interpolation.

## References

1. R. W. Hamming, *Numerical Methods for Scientist and Engineers*, 2<sup>nd</sup> edition, Dover Publications Inc., New York (1986).
2. W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in FORTRAN. The Art of Scientific Computing*, 2<sup>nd</sup> edition, Cambridge University Press (1994).



## Distribution

Admnstr  
Defns Techl Info Ctr  
Attn DTIC-OCP  
8725 John J Kingman Rd Ste 0944  
FT Belvoir VA 22060-6218

Central Intellgnc Agency  
Dir DB Standard  
Attn GE 47 QB (10 copies)  
Washington DC 20505

Defns Intllgnc Acgy  
Attn DT 2 Wpns & Sys Div  
Washington DC 20301

Dir of Defns Rsrch & Engrg  
Attn DD TWP  
Washintgon DC 20301

Commanding Officer  
Attn NMCB23  
6205 Stuart Rd Ste 101  
FT Belvoir VA 22060-5275

DARPA  
Attn Techl Lib  
3701 N Fairfax Dr  
Arlington VA 22203-1714

Dir of Chem & Nuc Ops DA DCSOPS  
Attn Techl Lib (2 copies)  
Washington DC 20310

Dpty Assist Scy for Rsrch & Techl  
Attn SARD-TR R Chait Rm 3E476  
Attn SARD-TT K Kominos  
Attn SARD-TT F Milton Rm 3E479  
Attn SARD-TT C Nash Rm 3E479  
The Pentagon  
Washington DC 20310-0103

Hdqtrs Dept of the Army  
Attn DAMO-FDQ D Schmidt  
Attn DAMO-FDQ MAJ M McGonagle  
400 Army Pentagon  
Washington DC 20310-0460

US Army Engrg Div  
Attn HNDED FD  
PO Box 1500  
Huntsville AL 35807

US Army Mis & Spc Intllgnc Ctr  
Attn AIAMS YDL  
Redstone Arsenal AL 35898-5500

US Army NGIC  
Attn Rsrch & Data Branch  
220 7th Stret NE  
Charlottesville VA 22901-5396

US Army Nuc & Cheml Agency  
7150 Heller Loop Ste 101 (3 copies)  
Springfield VA 22150-3198

US Army Rsrch Lab  
Attn SLCRO-D  
PO Box 12211  
Research Triangle Park NC 27709-2211

US Army Strtgc Defns Cmnd  
Attn CSSD H MPL Techl Lib  
Attn CSSD H XM Dr Davies  
PO Box 1500  
Huntsville AL 35807

Chief of Nav OPS Dept of the Navy  
Attn OP 03EG  
Attn OP 985F  
Washington DC 20350

Nav Civil Engrg Lab  
Attn Code L51 J Tancreto  
Port Hueneme CA 93043-5003

Officer in Charge  
Nav Construction Battalion Ctr  
Civil Engrg Lab  
Attn Code L31 Techl Lib  
Port Hueneme CA 93041

Nav Rsrch Lab  
Attn Code 2027 Techl Lib  
Washington DC 20375

## Distribution

Ofc of Nav Rsrch  
Attn Code 23 A Faulstick (2 copies)  
800 N Quincy Stret  
Arlington VA 22217

US Nav Academy  
Attn Tech Lib  
Attn P McCoy  
Attn LT D Fuller  
572 Holloway Rd  
Annapolis MD 21402-5002

US Nav Academy  
Attn R Smith  
363 Rickover Hall 590 Holloway Rd  
Annapolis MD 21402-5002

MIT  
Attn Tech Lib  
Cambridge MA 02139

Univ of Maryland  
Attn Engrg & Techl Lib  
Attn J Duncan  
Attn J Wallace  
Attn K Kreiger

Univ of Maryland (cont'd)  
Attn L Ong  
Attn P Bernard  
Rm 2168 Engrg Classroom Bldg  
College Park MD 20742-5121

US Army Rsrch Lab  
Attn AMSRL-CI-LL Tech Lib (3 copies)  
Attn AMSRL-CS-AL-TA Mail & Records  
Mgmt  
Attn AMSRL-CS-AL-TP Techl Pub  
(3 copies)  
Attn AMSRL-IS-EE A Wetmore  
Attn AMSRL-IS-EE D Garvey  
Attn AMSRL-IS-EE J Martin  
Attn AMSRL-IS-EE K Deacon  
Attn AMSRL-IS-EE R Cianco  
Attn AMSRL-IS-EE R Loucks (5 copies)  
Attn AMSRL-IS-EE R Meyers  
Attn AMSRL-IS-EE R Pinnick  
Attn AMSRL-IS-EE S Niles  
Attn AMSRL-IS-ES COL Price  
Attn AMSRL-IS-ES P Gillespie  
Adelphi MD 20783-1197

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1996		3. REPORT TYPE AND DATES COVERED Interim, from 1 to 31 Oct. 1996
4. TITLE AND SUBTITLE Chebyshev Polynomial Fit for Terrain Elevation			5. FUNDING NUMBERS DA PR: B53A PE: P61120	
6. AUTHOR(S) Richard B. Loucks				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory Attn: AMSRL-IS-EE 2800 Powder Mill Road Adelphi, MD 20783-1197			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TN-86	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory 2800 Powder Mill Road Adelphi, MD 20783-1197			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES AMS code: 611102.53A11 ARL PR: 7FEJ60				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) There is currently a desire to use Chebyshev polynomials to fit terrain elevation data. Such a fit would create a surface function that exactly fits the known elevations, and would describe an elevation at any point on that surface. This note questions the appropriateness of using Chebyshev polynomials for this purpose, as opposed to linear interpolation or use of a cubic spline. A set of elevations in one direction is used to illustrate a point that large transitions in elevation influence the coefficients in the polynomial fit and contribute spectral energy to points far from the transition area. It argues that a linear interpolation process, or a cubic spline interpolation, would be more appropriate.				
14. SUBJECT TERMS Chebyshev, terrain, fit			15. NUMBER OF PAGES 13	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	